




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Metabolically unhealthy individuals, either with obesity or not, have a higher risk of critical coronavirus disease 2019 outcomes than metabolically healthy individuals without obesity

Nam Hoon Kim, Kyeong Jin Kim, Jimi Choi, Sin Gon Kim



PII: S0026-0495(21)00194-3

DOI: <https://doi.org/10.1016/j.metabol.2021.154894>

Reference: YMETA 154894

To appear in: *Metabolism*

Received date: 23 July 2021

Revised date: 22 September 2021

Accepted date: 26 September 2021

Please cite this article as: N.H. Kim, K.J. Kim, J. Choi, et al., Metabolically unhealthy individuals, either with obesity or not, have a higher risk of critical coronavirus disease 2019 outcomes than metabolically healthy individuals without obesity, *Metabolism* (2021), <https://doi.org/10.1016/j.metabol.2021.154894>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Metabolically unhealthy individuals, either with obesity or not, have a higher risk of critical coronavirus disease 2019 outcomes than metabolically healthy individuals without obesity

Nam Hoon Kim, MD\*, Kyeong Jin Kim, MD\*, Jimi Choi, PhD, Sin Gon Kim, MD

Authors' Affiliation

Division of Endocrinology and Metabolism, Department of Internal Medicine, Korea University College of Medicine, Seoul, South Korea

\*Nam Hoon Kim and Kyeong Jin Kim equally contributed to this work.

Corresponding author: Sin Gon Kim, M.D., Ph.D.

Division of Endocrinology and Metabolism, Department of Internal Medicine, Korea University Anam Hospital, Korea University College of Medicine, 73 Goryeodae-ro, Seongbuk-gu, Seoul 02841, Republic of Korea

Tel.: +82-2-920-5890, Fax: +82-2-953-9355, E-mail: [k50367@korea.ac.kr](mailto:k50367@korea.ac.kr)

## Abstract

*Background:* This study aimed to determine the relative and independent contributions of impaired metabolic health and obesity to critical coronavirus disease 2019 (COVID-19).

*Methods:* We analyzed 4,069 COVID-19 patients between January and June 2020 in South Korea, classified into four groups according to metabolic health status and body mass index (BMI): metabolically healthy normal weight (MHNW), metabolically unhealthy normal weight (MUNW), metabolically healthy obesity (MHO), and metabolically unhealthy obesity (MUO). The primary outcome was a composite of intensive care unit (ICU) admission, invasive mechanical ventilation (IMV), extracorporeal membrane oxygenation (ECMO), and death. Multivariable Cox proportional hazard regression models were used to estimate the hazard ratio (HR) for the outcome.

*Results:* The incidence rate (per 100 person-months) of critical COVID-19 was the lowest in the MHNW group (0.90), followed by the MHO (1.64), MUNW (3.37), and MUO (3.37) groups. Compared with MHNW, a significantly increased risk of critical COVID-19 was observed in MUNW (HR, 1.41; 95% CI, 1.01–1.98) and MUO (HR, 1.77; 95% CI, 1.39–2.44) but not in MHO (HR, 1.48; 95% CI, 0.98–2.23). The risk of ICU admission or IMV/ECMO was increased only in MUO; however, the risk of death was significantly higher in MUNW and MUO. The risk of critical COVID-19 increased insignificantly by 2% per 1 kg/m<sup>2</sup> BMI increase but significantly by 13% per 1 metabolically unhealthy component increase, even after mutually adjusting for BMI and metabolic health status.

*Conclusions:* Metabolic health is more important to COVID-19 outcomes than obesity itself, suggesting that metabolic health status should be considered for a precise and tailored management of COVID-19 patients.

*Keywords:* COVID-19, metabolic health, metabolic syndrome, body mass index, adverse clinical outcome.

Journal Pre-proof

## 1. Introduction

Earlier observations revealed that patients with severe forms of COVID-19 were more obese than those with non-severe disease [1,2]. Multiple studies have indicated that obesity, which is generally represented by a high body mass index (BMI), is associated with the severity and mortality of COVID-19 [3–7]. However, obesity is a predisposing condition for metabolic derangement and cardiovascular diseases [8], which have also been recognized as important risk factors for adverse outcomes of COVID-19 [9,10]. Therefore, the observed association between obesity and critical COVID-19 might be biased by the effects of comorbidities. Previous studies have attempted to solve this problem by adjusting for comorbidities such as diabetes, hypertension, and dyslipidemia [11–14]; however, this might be insufficient to identify a direct relationship between obesity and COVID-19 outcomes.

Many differences in the determining characteristics of metabolic phenotypes have been observed between East Asians and other ethnic groups. For example, a substantial number of East Asian patients with type 2 diabetes were reported to be nonobese, although the prevalence of type 2 diabetes was similar to that in Caucasians [15]. This observation directed much interest to metabolically unhealthy without obesity phenotypes. However, metabolically healthy with obesity phenotypes have also been observed [16].

We hypothesized that metabolic health beyond obesity might be crucial in determining the risk of COVID-19 outcomes. The objective of this large retrospective study was to investigate the relative contribution of obesity and metabolic health status to critical outcomes of COVID-19 patients in a nationwide cohort in South Korea.

## 2. Materials and methods

### ***2.1. Data source and study population***

The Korean National Health Insurance Service (NHIS), which is the sole mandatory public medical insurance system for all citizens of South Korea, recently released the NHIS-COVID-19 cohort database, which included 8080 patients with COVID-19 confirmed with a SARS-CoV-2 PCR test between January 1, 2020, and June 4, 2020. The database incorporated past longitudinal information of the patients before the diagnosis of COVID-19, including demographic, medical, and pharmaceutical data from 2015 to 2020. The relevant information included medical history based on the International Classification of Disease 10th revision (ICD-10), hospitalization including ICU admission, drug prescriptions, medical procedures, and anthropometric and biochemical laboratory information including body weight, height, waist circumference, systolic and diastolic blood pressure, fasting plasma glucose, and lipid profile. In addition, this database was merged with death records managed by the Korean National Statistical Office. All patients were followed up until 4 months after the diagnosis of COVID-19 or until death. A more detailed protocol has been previously published [17]. We presented the STORBE checklist in Supplementary Table S1.

This study was approved by the Institutional Review Board of Korea University Anam Hospital (approval no. 2020AN0482). The requirement for informed consent was waived because all patient data were anonymized and de-identified.

### ***2.2. Study definition and outcomes***

This is a nationwide retrospective cohort study. Supplementary Figure S1 shows the diagram of the selection of the study patients. Among patients with laboratory confirmed COVID-19, those with any missing variables for the identification of metabolic status were

excluded. We divided these patients into four groups according to the obesity and metabolic health status: (i) metabolically healthy normal weight (MHNW), (ii) metabolically unhealthy normal weight (MUNW), (iii) metabolically healthy obesity (MHO), and (iv) metabolically unhealthy obesity (MUO). Obesity was defined as BMI  $\geq 25$  kg/m<sup>2</sup> according to the Asia-Pacific BMI criteria, calculated as weight divided by height squared (kg/m<sup>2</sup>) [18]. A metabolically unhealthy status was defined as the presence of three or more of the following factors: (i) fasting plasma glucose level  $\geq 100$  mg/dL or current use of glucose-lowering agents under the ICD-10 codes for diabetes mellitus (E10–E14), (ii) blood pressure  $\geq 130/85$  mmHg or use of antihypertensive agents under the ICD-10 codes for hypertension (I10–I15), (iii) serum triglyceride level  $\geq 150$  mg/dL or current use of lipid-lowering agents under the ICD-10 code for dyslipidemia (E78), (iv) high-density lipoprotein-cholesterol level  $< 40$  mg/dL in men or  $< 50$  mg/dL in women or current use of lipid-lowering agents under the ICD-10 code for dyslipidemia (E78), and (v) waist circumference  $> 90$  cm in men or  $\geq 85$  cm in women based on the International Diabetes Federation criteria for Asians [18,19].

The primary outcome was a composite of ICU admission, invasive mechanical ventilation (IMV), extracorporeal membrane oxygenation (ECMO), and death of any cause from the diagnosis of COVID-19 to the end of follow-up.

### 2.3. Statistical analyses

Continuous data are presented as mean  $\pm$  standard deviation (SD) values for normally distributed variables and as medians and interquartile ranges for nonnormally distributed variables. Categorical data are presented as frequencies and percentages. ANOVA and Pearson's  $\chi^2$  test were used to compare baseline characteristics among the four groups.

We calculated the incidence of the primary composite outcome by dividing the total number



of events by the total follow-up period (person-months). Kaplan–Meier analysis followed by the log-rank test was used to examine differences in the cumulative incidence of the composite primary outcome according to the metabolic health and obesity phenotypes. We also used multivariable Cox proportional hazard regression models to analyze the hazard ratios (HRs) for the study outcomes according to each metabolic health and obesity phenotype. Model 1 was adjusted for age and sex, and model 2 was adjusted for all possible confounders including age, sex, smoking, alcohol consumption, physical activity, socioeconomic status, previous history of chronic pulmonary disease including asthma and chronic obstructive pulmonary disease. As general obesity (represented by a high BMI) and metabolic health are closely associated with each other, we further investigated their independent roles in the development of critical COVID-19 by further adjusting for BMI and metabolically healthy status mutually. For handling the missing data, available-case analyses (also known as pairwise deletion) were used without additional data handling.

All reported  $p$ -values were two-sided, and statistical significance was set at  $p < 0.05$ . All statistical analyses were performed using SAS software (version 9.4; SAS Institute Inc., Cary, NC, USA).

### 3. Results

Of patients with laboratory confirmed COVID-19 from the original NHIS-COVID-19 cohort database ( $n=8080$ ), those with any missing variables ( $n=4011$ ) for the identification of metabolic status, including BMI, waist circumference, fasting plasma glucose, blood pressure, and serum levels of triglyceride and high-density lipoprotein-cholesterol, were excluded. Finally, a total of 4069 patients were selected.

### 3.1. Baseline patient characteristics

The mean ( $\pm$ SD) age of the patients was  $55.5 \pm 14.2$  years, and 2539 (62.4%) were female. The mean BMI was  $24.0 \text{ kg/m}^2$ . A total of 879 patients (21.6%) had a history of cardiovascular disease, 1310 (32.2%) had hypertension and 1013 (24.9%) had chronic pulmonary disease. The baseline characteristics of patients in the four groups according to metabolic health and obesity status are listed in **Table 1**. In general, patients in the metabolically unhealthy groups (MUNW and MUO) were older than those in the metabolically healthy groups (MHNW and MHO). The proportion of male patients was the lowest in MHNW, followed by MUNW, MUO, and MHO. Metabolic parameters, including systolic/diastolic blood pressure and serum glucose/triglyceride levels, were higher in the metabolically unhealthy groups than in the metabolically healthy groups. Metabolically unhealthy patients also had a higher prevalence of comorbidities, including hypertension, diabetes mellitus, dyslipidemia, cardiovascular disease, chronic kidney disease, and chronic pulmonary disease than metabolically healthy patients.

### 3.2. Severe COVID-19 outcomes according to metabolic health and obesity status

During about 4 months of follow-up, 289 (7.1%) critical COVID-19 outcomes occurred. The incidence rate (per 100 person-months) of the outcomes was higher in metabolically unhealthy patients (3.37 in both MUNW and MUO) than in metabolically healthy patients (0.90 in MHNW and 1.64 in MHO). **Table 2** shows the HRs for the composite and individual COVID-19 outcomes in the four groups. Compared with the MHNW group as the reference, all three other groups had significantly higher HRs for the composite of critical COVID-19 outcomes in an unadjusted model (1.79 [1.19–2.70], 3.58 [2.59–4.95], and 3.59 [2.63–4.91] in MHO, MUNW, and MUO, respectively), indicating that both obesity and impaired metabolic health are important risk factors for critical COVID-19. However, after adjusting

for confounding variables (in two different models), patients with a metabolically unhealthy status (MUNW and MUO groups) still had a significantly higher risk of critical COVID-19, but patients in the MHO group did not. In model 2, the risk of critical COVID-19 was 41% higher in the MUNW group and 77% higher in the MUO group than in the MHNW group. The MUNW and MUO groups also had a significantly higher risk of death than the MHNW group, although the risks of ICU admission and IMV or ECMO requirement were higher only in the MUO group. Kaplan–Meier curves showed a time-dependent risk of each outcome in the four groups (**Fig. 1**). All outcomes were determined to occur early, mostly within 30 days after the diagnosis of COVID-19.

### ***3.3. Severe COVID-19 outcomes according to BMI category and the number of metabolically unhealthy components***

In an unadjusted model, the risk of critical COVID-19 was increased by 2% per 1 kg/m<sup>2</sup> increase in BMI and by 13% for every 1 component increase in metabolically unhealthy parameters (**Table 3**). However, after reciprocal adjustment for each variable and other confounding factors (in model 2), only the number of metabolically unhealthy parameters was associated with the risk of critical COVID-19.

## **4. Discussion**

In this retrospective cohort study, we found that impaired metabolic health was a stronger predictor of critical COVID-19 than obesity itself in the Korean population. Notably, metabolically unhealthy individuals, either with obesity or not, had a higher risk of critical COVID-19 outcomes, including ICU admission and death, than metabolically healthy individuals without obesity. In contrast, metabolically healthy individuals with obesity were

not associated with critical COVID-19. In addition, the risk of COVID-19 outcomes linearly increased with increasing number of metabolically unhealthy components independently of BMI, whereas BMI did not distinguish patients at a risk of critical COVID-19.

Metabolically unhealthy traits, generally manifesting as metabolic syndrome, have been proposed as important risk factors for adverse outcomes of COVID-19. Although the exact mechanisms underlying this association remain unknown, immunomodulation (such as a hyperimmune response) and increased expression of ACE2 (which is the viral entry receptor of SARS-CoV-2) partly explain the association between metabolic syndrome and COVID-19 outcomes [20,21]. However, these mechanisms may also be applicable to obesity-related COVID-19 outcomes. Thus, further studies are required to elucidate this issue.

Obesity is a heterogeneous condition. For example, individuals with the same BMI can show various phenotypes in terms of adipocyte biology, fat distribution, and body composition [22]. One of the important factors determining metabolically healthy and unhealthy obesity is the amount or distribution of visceral adipose tissue (VAT). Previous studies found that Asian diabetic patients had more VAT than Caucasian diabetic patients with the same waist circumference, which can explain the higher susceptibility of lean Asians to type 2 diabetes [23]. Given the close association of VAT with insulin resistance, and further with metabolic syndrome, this parameter could be a determinant of the metabolic health status at a given BMI [24]. Several studies have indicated that VAT could be an important indicator of COVID-19 severity. In a small-sample study, VAT was an independent risk factor for severe COVID-19 outcomes, but not BMI [25]. Another study showed that high VAT ( $\geq 128.5 \text{ cm}^2$ ) was a crucial factor in predicting COVID-19 severity [26]. Although we were unable to validate this association in our cohort, we assumed that a higher proportion of VAT in the MUNW group than in the MHNW group might have contributed to the risk of critical COVID-19. Metabolically unhealthy phenotype is also related to adipose tissue dysfunction

and inflammation, which ultimately result in ectopic fat deposition and insulin resistance [27]. Indeed, upregulation of inflammatory cytokines including TNF- $\alpha$  and IL-6 was frequently observed in patients with insulin resistance, even without general obesity [20]. This partly explains hyperimmune response to infectious insult including SARS-CoV-2 and adverse outcomes of COVID-19 in MUNW people. Another possible factor is reduced muscle mass characteristic of MUNW individuals [16]. Reduced muscle mass has been proposed to be independently associated with insulin resistance and increased susceptibility to adverse respiratory outcomes, including pneumonia and sepsis [28,29].

Nonetheless, the available evidence from multiple geographical regions generally supports that obesity contributes to the adverse outcomes of COVID-19. However, previous reports had substantial inconsistencies in terms of study population and measured outcomes. A linear association between BMI and the need for ICU admission or IMV was reported in multiple studies [5,30,31]. However, for the mortality risk from COVID-19, J-shaped or U-shaped associations with increasing BMI were frequently observed, indicating that both severe obesity and underweight are risk factors for fatal events related to COVID-19 [5,7,30,32]. Given that the proposed cutoff values of high BMI for death risk ranged from 30 to 40 kg/m<sup>2</sup> in previous studies, the possibility exists that the much lower incidence of high BMI in our study population, with only 4.1% having a BMI of > 30 kg/m<sup>2</sup>, may not detect the harmful effects of severe obesity. We also found a J-shaped association between BMI and critical COVID-19, with patients having a BMI of between 20 and 24.9 kg/m<sup>2</sup> having the lowest risk of severe outcomes, although it was not significant. Therefore, ethnicity may be a crucial factor in determining the association between BMI and disease severity. Further, the fact that MUO patients still had the highest risk of severe COVID-19 and mortality in this study should not be ignored.

A notable result of our study was that the risk of critical COVID-19 increased with the

increasing number of metabolically unhealthy components. This was significant even after adjusting for BMI, suggesting an independent association between metabolic derangement, rather than obesity itself, and COVID-19 severity. We found that individuals with four or more metabolic syndrome components had a two-fold higher risk of COVID-19 outcomes than metabolically healthy individuals.

Taken together, impaired metabolic health, represented by insulin resistance, is closely related to the severity of COVID-19. In this regard, it is worth paying attention to the research results that several drugs related to improvement in insulin resistance, particularly metformin and peroxisome proliferator-activated receptors agonists, are associated with better outcomes in COVID-19 [33,34]

This study had strengths and limitations. The cohort database included anthropometric and laboratory measurements before the diagnosis of COVID-19, which made it possible to identify the underlying metabolic conditions of the patients. In addition, body weight, height, and waist circumference were measured using standardized methods (not self-reported). However, only 50% of the cohort had data on all analyzed metabolic syndrome components, resulting in the exclusion of a large proportion of patients. Our data also included detailed information on patient care, such as dates of hospital admission and ICU admission, medical procedures, and death records. Thus, it was possible to analyze the time-dependent association between exposure and outcomes. Although the follow-up duration was relatively short (up to 4 months), we found that most outcomes occurred within 2 months after the diagnosis of COVID-19, indicating that the follow-up duration would not limit the reliability of the study results. Finally, there might be residual confounding by unmeasured covariates, such as quality of intensive care, occupation, and viral load of SARS-CoV-2 in affected individuals.

In conclusion, our study showed that metabolically unhealthy traits are important risk

factors for critical COVID-19 independent of BMI in the Korean population. Considering that COVID-19 remains a tremendous public health problem, this study suggests that the metabolic health status of affected individuals should be evaluated and more attention needs to be paid to managing individuals with metabolically unhealthy traits.

### ***Acknowledgments***

We thank the participants of the Korean Health Insurance Cohort study and the National Health Insurance Service, which developed the NHIS-COVIL-19 cohort database.

### ***Funding***

This study was supported by a Korea University grant K1625581.

### ***Conflict-of-interest statement***

The authors report no potential conflicts of interest relevant to this article.

### ***Author contributions***

N.H.K. and K.J.K. contributed to data interpretation and wrote the first draft of the manuscript. J.C. performed the data analysis and contributed to data interpretation. S.G.K. designed the study and critically reviewed the manuscript. S.G.K. is the guarantor of this work, has full access to all study data, and takes responsibility for the integrity of the data and the accuracy of the data analysis.

## References

1. Cai Q, Chen F, Wang T, Luo F, Liu X, Wu Q, He Q, Wang Z, Liu Y, Liu L, Chen J, Xu L. Obesity and COVID-19 severity in a designated hospital in Shenzhen, China. *Diabetes Care* 2020;43:1392-1398
2. Bhatraju PK, Ghassemieh BJ, Nichols M, Kim R, Jerome KR, Nalla AK, Greninger AL, Pipavath S, Wurfel MM, Evans L, Kritek PA, West TE, Luks A, Gerbino A, Dale CR, Goldman JD, O'Mahony S, Mikacenic C. Covid-19 in critically ill patients in the Seattle region – case series. *N Engl J Med* 2020;382:2012-2022
3. Price-Haywood EG, Burton J, Fort D, Seoane L. Hospitalization and mortality among black patients and white patients with Covid-19. *N Engl J Med* 2020;382:2534-2543
4. Sattar N, Ho FK, Gill JM, Ghouri N, Gray SJ, Celis-Morales CA, Katikireddi SV, Berry C, Pell JP, McMurray JJ, Welsh P. BMI and future risk for COVID-19 infection and death across sex, age and ethnicity: preliminary findings from UK biobank. *Diabetes Metab Syndr* 2020;14:1149-1151
5. Gao M, Piernas C, Astbury NM, Hippisley-Cox J, O'Rahilly S, Aveyard P, Jebb SA. Associations between body-mass index and COVID-19 severity in 6.9 million people in England: a prospective, community-based, cohort study. *Lancet Diabetes Endocrinol* 2021;9:350-359
6. Palaiodimos L, Kokkinidis DG, Li W, Karamanis D, Ognibene J, Arora S, Southern WN, Mantzoros CS. Severe obesity, increasing age and male sex are independently associated with worse in-hospital outcomes, and higher in-hospital mortality, in a cohort of patients with COVID-19 in the Bronx, New York. *Metabolism* 2020;108:154262
7. Guerson-Gil A, Palaiodimos L, Assa A, Karamanis D, Kokkinidis D, Chamorro-Pareja N, Kishore P, Leider JM, Lawrence J Brandt. Sex-specific impact of severe obesity in the outcomes of hospitalized patients with COVID-19: a large retrospective study from the Bronx,



New York. *Eur J Clin Microbiol Infect Dis* 2021;40:1963-1974

8. Sowers JR. Obesity and cardiovascular disease. *Clin Chem* 1998;44:1821-1825

9. Xie J, Zu Y, Alkhatib A, Pham TT, Gill F, Jang A, Radosta S, Chaaya G, Myers L, Zifodya JS, Bojanowski CM, Marrouche NF, Mauvais-Jarvis F, Denson JL. Metabolic syndrome and COVID-19 mortality among adult black patients in New Orleans. *Diabetes Care* 2020;44:188-193

10. Lim S, Bae JH, Kwon HS, Nauck MA. COVID-19 and diabetes mellitus: from pathophysiology to clinical management. *Nat Rev Endocrinol* 2021;17:11-30

11. Gao F, Zheng KI, Wang XB, Sun QF, Pan KH, Wang TY, Chen YP, Targher G, Byrne CD, George J, Zheng MH. Obesity is a risk factor for greater COVID-19 severity. *Diabetes Care* 2020;43:e72-e74

12. Kim L, Garg S, O'Halloran A, Whitaker M, Pham H, Anderson EJ, Armistead I, Bennett NM, Billing L, Como-Sabetti K, Hill M, Kim S, Monroe ML, Muse A, Reingold AL, Schaffner W, Sutton M, Talbot HK, Torres SM, Yousey-Hindes K, Holstein R, Cummings C, Brammer L, Hall AJ, Fry AM, Lingley GE. Risk factors for intensive care unit admission and in-hospital mortality among hospitalized adults identified through the US Coronavirus Disease 2019 (COVID-19)-Associated Hospitalization Surveillance Network (COVID-NET). *Clin Infect Dis* 2021;72:e206-e214

13. Yates T, Zaccardi F, Islam N, Razieh C, Gillies CL, Lawson CA, Chudasama Y, Rowlands A, Davies MJ, Docherty AB, Openshaw PJM, Baillie JK, Semple MG, Khunti K. Obesity, ethnicity, and risk of critical care, mechanical ventilation, and mortality in patients admitted to hospital with COVID-19: analysis of the ISARIC CCP-UK Cohort. *Obesity (Silver Spring)* 2021;29:1223-1230

14. Chetboun M, Raverdy V, Labreuche J, Simonnet A, Wallet F, Caussy C, Antonelli M, Artigas A, Goma G, Meziani F, Helms J, Mylonakis E, Levy M, Kalligeros M, Latronico N,

- Piva S, Cerf C, Neuville M, Klouche K, Larchar R, Tamion F, Occhiali E, Snacken M, Preiser JC, Kontar L, Riviere A, Silva S, Sarton B, Krouchi R, Dubar V, Palaodimos L, Karamanis D, Perche J, L'Her E, Busetto L, Dicker D, Lev S, Duhamel A, Jourdain M, Pattou F. Association of body mass index and other metabolic risk factors with pneumonia outcomes in critically ill patients with coronavirus disease-19: an international multicenter retrospective cohort study. 2020. Available from <https://doi.org/10.2139/ssrn.3667634>
15. Rhee EJ. Diabetes in Asians. *Endocrinol Metab (Seoul)* 2015;30:263-269
  16. Stefan N. Metabolically healthy and unhealthy normal weight and obesity. *Endocrinol Metab (Seoul)* 2020;35:487-493
  17. Cho DH, Lee SJ, Jae SY, Kim WJ, Ha SJ, Gwon JG, Choi J, Kim DW, Kim JY. Physical activity and the risk of COVID-19 infection and mortality: a nationwide population-based case-control study. *J Clin Med* 2021;10
  18. The Asia-Pacific Perspective: Redefining Obesity and its Treatment [article online], 2000. Available from <https://apps.who.int/iris/handle/10665/206936>
  19. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JJ, Donato KA, Fruchart JC, James WP, Loria CM, Smith SC Jr. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;120:1640-1645
  20. Batabyal R, Freishtat N, Hill E, Rehman M, Freishtat R, Koutroulis I. Metabolic dysfunction and immunometabolism in COVID-19 pathophysiology and therapeutics. *Int J Obes (Lond)* 2021;45:1163-1169
  21. Stefan N, Birkenfeld AL, Schulze MB. Global pandemics interconnected – obesity, impaired metabolic health and COVID-19. *Nat Rev Endocrinol* 2021;17:135-149

22. Stefan N, Schick F, Häring HU. Causes, characteristics, and consequences of metabolically unhealthy normal weight in humans. *Cell Metab* 2017;26:292-300
23. Kadowaki T, Sekikawa A, Murata K, Maegawa H, Takamiya T, Okamura T, El-Saed A, Miyamatsu N, Edmundowicz D, Kita Y, Sutton-Tyrrell K, Kuller LH, Ueshima H. Japanese men have larger areas of visceral adipose tissue than Caucasian men in the same levels of waist circumference in a population-based study. *Int J Obes (Lond)* 2006;30:1163-1165
24. Hayashi T, Boyko EJ, McNeely MJ, Leonetti DL, Kahn SE, Fujimoto WY. Visceral adiposity, not abdominal subcutaneous fat area, is associated with an increase in future insulin resistance in Japanese Americans. *Diabetes* 2008;57:1269-1275
25. Petersen A, Bressan K, Albrecht J, Thieß HM, Vaidiek J, Hamm B, Makowski MR, Niehues A, Niehues SM, Adams LC. The role of visceral adiposity in the severity of COVID-19: highlights from a unicenter cross sectional pilot study in Germany. *Metabolism* 2020;110:154317
26. Favre G, Legueult K, Pradier C, Raffaelli C, Ichai C, Iannelli A, Redheuil A, Lucidarme O, Esnault V. Visceral fat is associated to the severity of COVID-19. *Metabolism* 2021;115:154440
27. Iacobini C, Pugliese G, Fantauzzi CB, Federici M, Menini S. Metabolically healthy versus metabolically unhealthy obesity. *Metabolism* 2019;92:51-60
28. Lee SW, Youm Y, Lee WJ, Choi W, Chu SH, Park YR, Kim HC. Appendicular skeletal muscle mass and insulin resistance in an elderly Korean population: the Korean social life, health and aging project-health examination cohort. *Diabetes Metab J* 2015;39:37-45
29. Okazaki T, Ebihara S, Mori T, Izumi S, Ebihara T. Association between sarcopenia and pneumonia in older people. *Geriatr Gerontol Int* 2020;20:7-13
30. Tartof SY, Qian L, Hong V, Wei R, Nadjafi RF, Fischer H, Li Z, Shaw SF, Caparosa SL, Nau CL, Saxena T, Rieg GK, Ackerson BK, Sharp AL, Skarbinski J, Naik TK, Murali SB.

Obesity and mortality among patients diagnosed with COVID-19: results from an integrated health care organization. *Ann Intern Med* 2020;173:773-781

31. Wu X, Li C, Chen S, Zhang X, Wang F, Shi T, Li Q, Lin L. Association of body mass index with severity and mortality of COVID-19 pneumonia: a two-center, retrospective cohort study from Wuhan, China. *Aging (Albany NY)* 2021;13:7767-7780

32. Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y, Tobin KA, Cerfolio RJ, Francois F, Horwitz LI. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ* 2020;369:m1966

33. Khunti K, Knighton P, Zaccardi F, Bakhai C, Burton E, Holman N, Kar P, Meace C, Sattar N, Sharp S, Wareham NJ, Weaver A, Welch E, Young B, Valabhji J. Prescription of glucose-lowering therapies and risk of COVID-19 mortality in people with type 2 diabetes: a nationwide observational study in England. *Lancet Diabetes Endocrinol* 2021;9:293-303

34. Yang W, Sun X, Zhang J, Zhang K. The effect of metformin on mortality and severity in COVID-19 patients with diabetes mellitus. *Diabetes Res Clin Pract* 2021;178:108977

**Table 1.** Characteristics of patients with coronavirus disease 2019 according to metabolic health and obesity phenotypes

	MHNW	MUNW	MHO	MUO	<i>p</i> -Value
n (%)	1871 (46.0)	728 (17.9)	595 (14.6)	875 (21.5)	
Follow-up time, mean (SD), days	118.0 (10.2)	111.9 (26.2)	117.0 (14.5)	113.0 (24.3)	
Age group, n (%), years					< 0.001
20–29	149 (8.0)	3 (0.4)	39 (6.6)	7 (0.8)	
30–39	228 (12.2)	9 (1.2)	40 (6.7)	41 (4.7)	
40–49	430 (23.0)	45 (6.2)	124 (20.8)	112 (12.8)	
50–59	604 (32.3)	179 (24.6)	192 (32.3)	238 (27.2)	
60–69	326 (17.4)	246 (33.6)	96 (16.1)	276 (31.5)	
70–79	95 (5.1)	165 (22.7)	41 (6.9)	135 (15.4)	
≥ 80	39 (2.1)	51 (11.1)	13 (2.2)	66 (7.5)	
Male sex, n (%)	522 (27.9)	300 (41.2)	298 (50.1)	410 (46.9)	< 0.001
BMI, mean (SD), kg/m <sup>2</sup>	21.8 (2.0)	22.7 (1.7)	27.0 (2.1)	27.9 (2.6)	< 0.001
WC, mean (SD), cm	74.6 (5.8)	80.2 (6.5)	85.8 (7.3)	90.7 (7.4)	< 0.001
Men	79.6 (6.1)	83.3 (5.5)	88.8 (6.6)	92.9 (7.2)	< 0.001
Women	72.7 (6.0)	78.0 (6.3)	82.8 (6.8)	88.7 (7.1)	< 0.001
SBP, mean (SD), mmHg	115.0 (13.2)	128.2 (15.2)	121.0 (13.1)	130.3 (14.4)	< 0.001
DBP, mean (SD), mmHg	71.4 (9.2)	77.6 (9.9)	75.3 (8.9)	79.7 (9.9)	< 0.001
FPG, mean (SD), mg/dL	93.4 (15.9)	113.3 (34.7)	93.9 (12.1)	113.6 (31.1)	0.001
Total cholesterol, mean (SD), mg/dL	192.0 (34.7)	196.6 (44.5)	197.1 (31.7)	197.3 (43.0)	< 0.001
Triglyceride, mean (SD), mg/dL	92.1 (47.3)	150.2 (91.1)	111.8 (60.3)	165.5 (119.2)	< 0.001
HDL-C, mean (SD), mg/dL	61.4 (17.4)	53.5 (14.4)	55.5 (12.8)	50.4 (12.3)	< 0.001
Hemoglobin, mean (SD), g/dL	13.4 (1.5)	13.7 (1.5)	14.1 (1.6)	14.2 (1.6)	< 0.001
Creatinine, mean (SD), mg/dL	0.80 (0.29)	0.86 (0.30)	0.86 (0.20)	0.90 (0.38)	< 0.001

AST, mean (SD), IU/L	22.5 (8.6)	26.2 (13.2)	25.4 (19.0)	31.1 (37.0)	< 0.001
ALT, mean (SD), IU/L	19.0 (12.1)	23.8 (13.2)	25.6 (25.5)	34.8 (52.5)	< 0.001
Smoking, n (%)					< 0.001
Never	1577 (84.3)	500 (68.7)	444 (74.6)	612 (69.9)	
Former	200 (10.7)	120 (16.5)	100 (16.8)	184 (21.0)	
Current	94 (5.0)	58 (8.0)	51 (8.6)	79 (9.0)	
Alcohol consumption, n (%)	313 (16.7)	106 (14.6)	159 (26.7)	203 (23.2)	< 0.001
Regular exercise, n (%)	805 (43.0)	338 (46.4)	262 (44.0)	429 (49.0)	0.023
Moderate-to-vigorous physical activity, n (%)	93 (5.0)	45 (6.2)	52 (8.4)	45 (5.1)	0.663
Low SES*, n (%)	430 (23.0)	178 (24.5)	117 (19.7)	199 (22.7)	0.214
Comorbidities, n (%)					
Hypertension	200 (10.7)	464 (63.7)	95 (16.0)	551 (63.0)	< 0.001
Diabetes mellitus	33 (1.8)	254 (36.3)	8 (1.3)	287 (32.8)	< 0.001
Dyslipidemia	162 (8.7)	597 (82.0)	41 (6.9)	617 (70.5)	< 0.001
Cardiovascular disease	227 (12.1)	294 (40.4)	66 (11.1)	292 (33.4)	< 0.001
Chronic kidney disease	8 (0.4)	22 (3.0)	4 (0.7)	35 (4.0)	< .0001
Chronic pulmonary disease	324 (20.5)	227 (31.2)	121 (20.3)	281 (32.1)	< 0.001

Abbreviations: MHNW, metabolically healthy normal weight; MUNW, metabolically unhealthy normal weight; MHO, metabolically healthy obesity; MUO, metabolically unhealthy obesity; SD, standard deviation; BMI, body mass index; WC, waist circumference; SBP, systolic blood pressure; DBP, diastolic blood pressure; FPG, fasting plasma glucose; HDL-C, high-density lipoprotein-cholesterol; AST, aspartate aminotransferase; ALT, alanine aminotransferase; SES, socioeconomic status.

\*Socioeconomic status was identified according to the medical insurance premium in the database, in which low socioeconomic status was defined as the lower 30%.

**Table 2.** Severe COVID-19 outcomes according to metabolic health and obesity phenotypes

	Events, n	Incidence rate, per 100 person- months	Hazard ratio (95% CI)		
			Unadjusted	Model 1	Model 2
<i>Composite of severe COVID-19 outcomes</i>					
MHNW	64	0.90	1 (ref.)	1 (ref.)	1 (ref.)
MUNW	86	3.37	3.58 (2.59–4.95)	1.42 (1.02–1.97)	1.41 (1.01–1.98)
MHO	36	1.64	1.79 (1.19–2.70)	1.42 (0.94–2.14)	1.48 (0.98–2.23)
MUO	103	3.37	3.59 (2.63–4.91)	1.30 (1.31–2.47)	1.77 (1.29–2.44)
<i>ICU admission</i>					
MHNW	47	0.66	1 (ref.)	1 (ref.)	1 (ref.)
MUNW	50	1.96	2.82 (1.89–4.40)	1.43 (0.94–2.18)	1.45 (0.95–2.20)
MHO	29	1.32	1.96 (1.24–3.12)	1.60 (1.00–2.55)	1.59 (0.99–2.54)
MUO	69	2.26	3.26 (2.25–4.72)	1.92 (1.31–2.81)	1.92 (1.31–2.82)
<i>IMV or ECMO</i>					
MHNW	15	0.21	1 (ref.)	1 (ref.)	1 (ref.)
MUNW	24	0.91	4.19 (2.20–7.98)	1.83 (0.94–3.58)	1.87 (0.95–3.65)
MHO	12	0.53	2.54 (1.19–5.42)	1.99 (0.93–4.27)	2.02 (0.94–4.33)
MUO	35	1.10	5.11 (2.79–9.36)	2.72 (1.47–5.05)	2.65 (1.42–4.94)
<i>Death</i>					
MHNW	21	0.29	1 (ref.)	1 (ref.)	1 (ref.)
MUNW	52	1.94	6.56 (3.95–10.89)	1.85 (1.11–3.10)	1.90 (1.13–3.19)
MHO	12	0.52	1.81 (0.89–3.67)	1.30 (0.64–2.65)	1.44 (0.71–2.94)
MUO	52	1.60	5.43 (3.27–9.01)	2.23 (1.34–3.71)	2.22 (1.33–3.70)

Abbreviations: COVID-19, coronavirus disease 2019; MHNW, metabolically healthy normal weight; MUNW, metabolically unhealthy normal weight; MHO, metabolically healthy obesity; MUO, metabolically unhealthy obesity; ICU, intensive care unit; IMV, invasive mechanical ventilation; ECMO, extracorporeal membrane

oxygenation.

Model 1 was adjusted for age and sex.

Model 2 was adjusted for age, sex, smoking, alcohol consumption, physical activity, socioeconomic status, and previous history of chronic pulmonary disease, including asthma and chronic obstructive pulmonary disease.



**Table 3.** Severe coronavirus disease 2019 outcomes according to body mass index category and the number of metabolically unhealthy components

	n	Events, n	Incidence rate, per 100 person- months	Hazard ratio (95% CI)		
				Unadjusted	Model 1	Model 2
<i>BMI, kg/m<sup>2</sup></i>						
< 20	393	25	1.72	1.07 (0.68–1.69)	1.22 (0.77–1.92)	1.22 (0.77–1.93)
20–22.9	1194	71	1.60	1 (ref.)	1 (ref.)	1 (ref.)
23–24.9	1012	54	1.43	0.90 (0.63–1.28)	0.78 (0.55–1.11)	0.72 (0.51–1.03)
25–29.9	1279	126	2.77	1.69 (1.27–2.27)	1.28 (0.96–1.72)	1.19 (0.88–1.60)
≥ 30	191	13	1.86	1.15 (0.64–2.08)	1.46 (0.81–2.64)	1.16 (0.63–2.14)
<i>p-Value for trend</i>				0.002	0.129	0.118
Per 1 kg/m <sup>2</sup> increase				1.04 (1.01–1.06)	1.03 (0.99–1.07)	1.02 (0.98–1.06)
<i>No. of MU components</i>						
0	945	19	0.52	1 (ref.)	1 (ref.)	1 (ref.)
1	793	35	1.17	2.22 (1.27–3.88)	1.41 (0.81–2.48)	1.14 (0.80–2.49)
2	728	46	1.71	3.21 (1.88–5.48)	1.47 (0.85–2.52)	1.43 (0.83–2.49)
3	650	55	2.25	4.34 (2.58–7.32)	1.65 (0.97–2.81)	1.62 (0.94–2.80)
4	649	91	4.09	7.36 (4.49–12.06)	2.14 (1.28–3.56)	2.04 (1.20–3.47)
5	304	43	4.16	7.47 (4.35–12.82)	2.02 (1.16–3.54)	1.84 (0.99–3.41)
<i>p-Value for trend</i>				< 0.001	0.001	0.001
Per 1 component increase				1.47 (1.37–1.59)	1.15 (1.06–1.25)	1.13 (1.03–1.24)

Abbreviations: BMI, body mass index; MU, metabolically unhealthy; ref., reference.

Model 1 was adjusted for age and sex.

Model 2 was adjusted for age, sex, smoking, alcohol consumption, physical activity, socioeconomic status, and previous history of chronic pulmonary disease (including asthma and chronic obstructive pulmonary disease) and further adjusted for metabolically unhealthy status and BMI.

## Figure Legends

Figure 1. Kaplan–Meier analysis of severe coronavirus disease 2019 outcomes (A), ICU admission (B), IMV or ECMO therapy (C), and death (D) according to metabolic health and obesity phenotypes.

\*Abbreviations: MHNW, metabolically healthy weight; MUNW, metabolically unhealthy normal weight; MHO, metabolically healthy obesity; MUO, metabolically unhealthy obesity; ICU, intensive care unit; IMV, invasive mechanical ventilation; ECMO, extracorporeal membrane oxygenation.

*CRedit authorship contribution statement*

**Nam Hoon Kim:** Data curation, Investigation, Writing-Original draft & review and editing.

**Kyeong Jin Kim:** Data curation, Investigation, Writing-Original draft.

**Jimi Choi:** Methodology, Formal analysis, Data curation.

**Sin Gon Kim:** Conceptualization, Investigation, Supervision, Project administration.

Journal Pre-proof

**Highlight**

- Analyzing a nation-wide COVID-19 cohort database.
- The risk of death was higher in metabolic unhealthy normal and obesity patients.
- Metabolic health is more important to COVID-19 outcomes than obesity.







Figure 1